

RAYLEIGH-TAYLOR STABILITY OF A STRONG VERTICAL MAGNETIC FIELD AT THE GALACTIC CENTER CONFINED BY A DISK THREADED WITH HORIZONTAL MAGNETIC FIELD

B. D. G. CHANDRAN^a

DEPT. PHYSICS & ASTRONOMY, UNIVERSITY OF IOWA, IOWA CITY, IA

^a E-MAIL: BENJAMIN-CHANDRAN@UIOWA.EDU

submitted to ApJ

ABSTRACT

Observations of narrow radio-emitting filaments near the Galactic center have been interpreted in previous studies as evidence of a pervasive vertical (i.e. $\perp$ to the Galactic plane) milliGauss magnetic field in the central ~ 150 pc of the Galaxy. A simple cylindrically symmetric model for the equilibrium in this central region is proposed in which horizontal (i.e. $\parallel$ to the Galactic plane) magnetic fields embedded in an annular band of partially ionized molecular material of radius ~ 150 pc are wrapped around vertical magnetic fields threading low-density hot plasma. The central vertical magnetic field, which has a pressure that significantly exceeds the thermal pressure of the medium, is confined by the weight of the molecular material. The stability of this equilibrium is studied indirectly by analyzing a uniformly rotating cylinder of infinite extent along the z axis in cylindrical coordinates (r, θ, z) , with low-density plasma and an axial magnetic field at $r < 150$ pc, high-density plasma and an azimuthal field at $r > 150$ pc, and a gravitational acceleration $g^* \propto r$ directed in the $-\hat{r}$ direction. Simple profiles are assumed for the density ρ , pressure p , and field strength B , with the sound speed and Alfvén speeds $\propto r$ within the dense plasma. The density profile and gravity tend to destabilize the plasma, but the plasma tends to be stabilized by rotation and magnetic tension—since the interface between the high and low-density plasmas can not be perturbed without bending either the horizontal or vertical field. Normal modes proportional to $e^{(im\theta + ik_z z - i\sigma t)}$ with $k_z = 0$ and $m \neq 0$ are studied. Such modes neither bend nor compress the axial field at $r < 150$ pc but allow compressions of the dense plasma along the azimuthal magnetic field that enhance the destabilizing role of gravity. It is shown analytically that when $\beta = 8\pi p/B^2$ is small and the dense plasma is supported against gravity primarily by rotation, the necessary and sufficient condition for stability to $k_z = 0$ modes is $|g| < 2|\Omega|a$, where $g = g^* - \Omega^2 r$ is the effective gravity, Ω is the uniform angular velocity, and a is the sound speed in the dense plasma. Since the effective gravity is determined by the degree to which magnetic (and to a lesser degree pressure) forces support the dense plasma, the stability criterion gives an upper limit on the strength of the axial magnetic field, which is ~ 1 mG for Galactic-center parameters.

1. INTRODUCTION

The central ~ 150 pc of the Galaxy host a phenomenon that is unique within the Galaxy: narrow, “vertical” (i.e. $\perp$ to the Galactic plane) filaments of radio emission that are typically a few tens of parsecs long and only a fraction of a parsec wide (Morris 1996). Polarization measurements indicate that the magnetic fields in the Radio Arc and Northern Thread are parallel to these filaments (Tsuboi et al. 1986, Reich 1994, Tsuboi et al. 1995, Lang et al. 1999b), and this alignment is presumed to be typical of the other filaments as well. In virtually all cases the filaments appear to be in contact with a molecular cloud at some point along their length. The absence of bending of these filaments by the randomly moving clouds has been interpreted as evidence of a milliGauss lower limit to the field strength in the filaments (Yusef-Zadeh & Morris 1987a—this point is discussed further in section 7). The apparent impossibility of forming or confining narrow filaments with such tremendous magnetic pressures—well in excess of the thermal pressure of the medium—suggests that if a mG vertical field is present, it pervades the central ~ 150 pc of the Galaxy, and the filaments are simply those flux tubes that possess large populations of relativistic electrons (Morris & Serabyn 1996).

A strong pervasive magnetic field at the Galactic center would be a natural consequence of radial inflow in the Galactic disk (Sofue & Fujimoto 1987, Morris & Serabyn 1996), and magnetic-pressure forces would over time cause the volume-

filling magnetic flux tubes to rise buoyantly away from the Galactic mid-plane, forcing field lines that penetrate the mid-plane into a vertical orientation (Chandran et al. 2000). It is not clear, however, that such a scenario is inevitable, as the rate of radial inflow in the Galactic disk is not well known and since sufficient turbulent resistivity can inhibit the central concentration of the field (Lubow, Papaloizou, & Pringle 1994). Alternative explanations of the filaments have been explored by a number of authors (Chudnovsky et al. 1986, Benford 1988, Heyvaerts, Norman, & Pudritz 1988, Lesch & Reich 1992, Rosso & Pelletier 1993, Shore & LaRosa 1999). If a pervasive mG field is present, it would play a dominant role in the dynamics of the interstellar medium near the Galactic center, and would suggest that the Galaxy was born with a magnetic field $\gtrsim 10^{-7}$ G (Chandran et al 2000—this point is discussed further in section 8).

One of the most pressing questions regarding the pervasive-mG-field hypothesis is whether such a strong magnetic field could be stably confined. The volume-dominant phase of the central region is hot plasma at a temperature $T \sim 10^8$ K which fills an elliptical region 150×270 pc (FWHM) with a major axis that is tilted 20° with respect to the Galactic plane (Yamauchi et al. 1990). The density of this plasma was estimated to be $0.03 - 0.06 \text{ cm}^{-3}$ by Yamauchi et al. (1990) and $\sim 0.3 - 0.4$ by Koyama et al. (1996). The mass-dominant phase of the central region is molecular gas with $n > 10^4 \text{ cm}^{-3}$, $T \sim 70$ K, a

filling factor in excess of 0.1, and a vertical thickness of ~ 30 pc (Bally et al. 1988, Morris & Serabyn 1996). If the magnetic field strength is 1 mG, then the magnetic pressure $B^2/8\pi$ is $\sim 4 \times 10^{-8}$ dyne/cm², which far exceeds the thermal pressure of both the hot plasma, $p_{\text{hot}} \sim 4 \times 10^{-10} - 4 \times 10^{-9}$ dyne/cm², and the molecular material, $p_{\text{cloud}} \sim 10^{-10}$ dyne/cm². The thermal pressure of the ambient medium is thus unable to confine mG fields, whether they are ubiquitous or in the form of isolated narrow flux tubes.

The only force capable of confining a pervasive vertical mG field is the weight of the molecular material. Interestingly, the polarization of dust emission indicates that the molecular material is threaded by horizontal (i.e., $\parallel$ to the Galactic plane) magnetic fields (Morris & Serabyn 1996). Because this horizontal field is perpendicular to the vertical magnetic field in the hot plasma, it is difficult for the two types of material to interpenetrate, which suggests that the weight of the molecular material may be able to stably prevent a strong vertical field from expanding. Although the molecular clouds are clumpy, it is expected that the horizontal magnetic field completely encloses the central region. In this paper, the equilibrium of figure 1 is proposed as a starting point for investigating the confinement of vertical fields at the Galactic center.

To make a compact analytic treatment possible, the stability of this equilibrium is explored under a number of additional simplifying approximations. It is assumed that the fractional ionization within the molecular clouds is sufficiently large and the frequencies of any instabilities sufficiently small compared to the neutral-ion collision frequency that ambipolar diffusion can be ignored. The partially ionized clouds then behave as a single-species fluid to which the magnetic field is frozen, or, in other words, as a dense plasma described by the equations of magnetohydrodynamics (MHD). The equilibrium is taken to be infinite and invariant along the spin axis—the z axis in cylindrical coordinates (r, θ, z) —as depicted in figure 2. A constant angular velocity Ω is assumed, which removes the magneto-rotational instability from the problem (Balbus & Hawley 1998). An axial magnetic field and low-density plasma are assumed for $r < 150$ pc, and an azimuthal magnetic field and dense plasma for $r > 150$ pc. The equilibrium density ρ_0 , pressure p_0 , and field strength B_0 are assumed to be proportional to powers of r within the dense plasma, and $\beta = 8\pi p_0/B_0^2$ is taken to be constant and small, which implies that the sound speed a and Alfvén speed v_A are proportional to r . The gravitational acceleration g^* is taken to be proportional to r and to be directed in the $-\hat{r}$ direction. If g^* arises from stars, this amounts to assuming a constant mass density of stars throughout the region occupied by the plasma. It is also assumed that the plasma is supported primarily by rotation, so that $v_A \ll v_{\text{rot}}$ in the dense plasma, where $v_{\text{rot}} = \Omega r$. Although some of these assumptions are extreme, a number of the key ingredients are retained: compressibility, the Coriolis force, the suspension of dense plasma “above” low-density plasma, and the orthogonal orientations of the magnetic fields in the low- and high-density plasmas.

The analysis then focuses on modes which perturb the sharp boundary between the high- and low-density plasmas. Such perturbations are to some extent stabilized by magnetic tension, since short-wavelength disturbances involve significant field-line bending of either the axial or azimuthal magnetic fields. Uniform rotation also plays a stabilizing role (e.g., Talwar 1960, Chandrasekhar 1961, Gilman 1970, Acheson & Gibbons 1978). Normal modes of the form $\mathbf{u}_1(r)e^{(ik_z z + im\theta - i\sigma t)}$ with $k_z = 0$ and

$m \neq 0$ are studied analytically. Such modes avoid bending or compressing the axial magnetic field at $r < 150$ pc and also allow compressions of the dense plasma along the azimuthal magnetic field that enhance the destabilizing effects of gravity. It is shown in sections 2 through 4 that the necessary and sufficient condition for stability to $k_z = 0$ modes is

$$|g| < 2|\Omega|a \quad (1)$$

when $a \ll v_A \ll \Omega r$, where $g = g^* - \Omega^2 r$ is the effective gravity, and a and v_A are evaluated in the dense plasma. A physical explanation of equation (1) is given in section 5. Since the effective gravity is proportional to the partial support of the plasma by the magnetic field, equation (1) can be re-written as an upper limit on the strength of the vertical field [equation (94)]. For Galactic-center parameters, the upper limit is ~ 1 mG. If equation (1) is the approximate criterion for the stability of the boundary between the high and low-density plasmas in systems with differential rotation, a finite extent along the z axis, and more general radial profiles—say by evaluating the terms in equation (1) near the inner edge of the dense plasma—then a pervasive mG vertical magnetic field can be confined at the Galactic center. Equation (1) would then also be relevant to other accreting systems with strong poloidal magnetic fields.

Global instabilities with $k_z \neq 0$, local quasi-interchange modes, and instabilities of the two-dimensional plasma of figure 1 are discussed in section 6. In section 7, observational constraints on the field-strength in the Galactic-center filaments are reviewed. The implications of a pervasive mG field at the Galactic center for the origin of the Galactic magnetic field are discussed in section 8.

2. DIFFERENTIAL EQUATION FOR $k_z = 0$ NORMAL MODES IN THE DENSE PLASMA

The plasma is taken to obey the equations of ideal MHD and an adiabatic equation of state,

$$\frac{\partial \rho}{\partial t} = -\nabla \cdot (\rho \mathbf{u}), \quad (2)$$

$$\rho \left(\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) = -\nabla \left(p + \frac{B^2}{8\pi} \right) + \frac{1}{4\pi} \mathbf{B} \cdot \nabla \mathbf{B} - \rho g^* \hat{r}, \quad (3)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{u} \times \mathbf{B}), \quad \text{and} \quad (4)$$

$$\left(\frac{\partial}{\partial t} + \mathbf{u} \cdot \nabla \right) \left(\frac{p}{\rho^\gamma} \right) = 0, \quad (5)$$

where ρ , p , $\mathbf{u}$, and $\mathbf{B}$ are the density, pressure, velocity, and magnetic field, g^* is the gravitational acceleration, (r, θ, z) are cylindrical coordinates, and γ is the adiabatic index. An equilibrium is assumed in which all quantities are functions of r alone. Equilibrium quantities are denoted with the subscript 0;

$$\rho_0 = \rho_0(r), \quad (6)$$

$$p_0 = p_0(r), \quad (7)$$

$$\mathbf{u}_0 = \Omega r \hat{\theta}, \quad \text{and} \quad (8)$$

$$\mathbf{B}_0 = B_0(r) \hat{\theta}. \quad (9)$$

A uniform angular velocity is assumed,

$$\Omega = \text{constant}, \quad (10)$$

and the effective gravity g is defined by the equation

$$g = g^* - \Omega^2 r. \quad (11)$$

By virtue of equation (3),

$$0 = \frac{d}{dr} \left(p_0 + \frac{B_0^2}{8\pi} \right) + \frac{B_0^2}{4\pi r} + \rho_0 g. \quad (12)$$

The sound and Alfvén speeds are defined, respectively, by the equations

$$a^2 = \frac{\gamma p_0}{\rho_0}, \quad (13)$$

and

$$v_A^2 = \frac{B_0^2}{4\pi\rho_0}. \quad (14)$$

Small-amplitude disturbances to the equilibrium, denoted by the subscript 1, can be decomposed into normal modes of the form

$$\mathbf{u}_1 = \mathbf{u}_1(r) e^{im\theta + ik_z z - i\sigma t}, \quad (15)$$

with analogous equations for $\mathbf{B}_1$, ρ_1 , and p_1 . The equations governing these disturbances are obtained in the standard way, by substituting $\mathbf{u} = \mathbf{u}_0 + \mathbf{u}_1$, $\mathbf{B} = \mathbf{B}_0 + \mathbf{B}_1$, etc into equations (2) through (5) and discarding all terms containing the product of more than one of the perturbed quantities. The analysis is confined to the case

$$k_z = 0. \quad (16)$$

The frequency ω of the disturbance in the frame rotating with the plasma is given by

$$\omega = \sigma - m\Omega. \quad (17)$$

From equations (2) and (5), the $\hat{r}$ and $\hat{\theta}$ components of equation (4), and the $\hat{\theta}$ component of equation (3), one obtains

$$B_{1r} = -\frac{mB_0 u_{1r}}{\omega r}, \quad (18)$$

$$B_{1\theta} = -\frac{i}{\omega} \frac{d}{dr} (u_{1r} B_0), \quad (19)$$

$$u_{1\theta} = -\frac{i}{D_s} \left(h u_{1r} + \frac{m a^2}{r} \frac{d u_{1r}}{dr} \right), \quad (20)$$

$$\rho_1 = \frac{-i u_{1r}}{\omega} \left[\frac{\rho_0}{r} \left(1 + \frac{mh}{D_s} \right) + \frac{d\rho_0}{dr} \right] - \frac{i\omega\rho_0}{D_s} \frac{d u_{1r}}{dr}, \quad (21)$$

and

$$p_1 = \frac{-i u_{1r}}{\omega} \left[\frac{a^2 \rho_0}{r} \left(1 + \frac{mh}{D_s} \right) + \frac{d p_0}{dr} \right] - \frac{i\omega\rho_0 a^2}{D_s} \frac{d u_{1r}}{dr}, \quad (22)$$

where

$$h = 2\Omega\omega - \frac{m}{r} \left(g - \frac{a^2}{r} \right), \quad (23)$$

and

$$D_s = \omega^2 - \frac{m^2 a^2}{r^2}. \quad (24)$$

Upon substituting equations (18) through (22) into the $\hat{r}$ component of equation (3), one obtains

$$\frac{d}{dr} \left[\rho_0 \left(v_A^2 + \frac{a^2 \omega^2}{D_s} \right) r \frac{d u_{1r}}{dr} \right] + r F u_{1r} = 0, \quad (25)$$

where

$$F = \rho_0 D_A - \frac{\rho_0 h^2}{D_s} + \frac{1}{r} \frac{d}{dr} \left(\frac{\rho_0 a^2 m h}{D_s} \right) + \frac{d}{dr} \left(\frac{\rho_0 a^2}{r} \right) + \frac{B_0^2}{4\pi r^2} - \rho_0 r \frac{d}{dr} \left(\frac{g}{r} \right), \quad (26)$$

and where

$$D_A = \omega^2 - \frac{m^2 v_A^2}{r^2}. \quad (27)$$

For isothermal equilibria and $\gamma = 1$, equation (25) reduces to the $k_z = 0$ limit of equation (2.13) of Acheson & Gibbons (1978).

Because $k_z = 0$, u_{1z} and B_{1z} decouple from the other variables and exhibit purely stable behavior: the $\hat{z}$ component of equation (4) gives

$$B_{1z} = -\frac{m B_0 u_{1z}}{\omega r}, \quad (28)$$

and the $\hat{z}$ component of equation (3) then gives

$$D_A u_{1z} = 0. \quad (29)$$

3. DIFFERENTIAL EQUATION FOR $k_z = 0$ NORMAL MODES IN THE CONSTANT- β EQUILIBRIUM

It is now assumed that the equilibrium has constant β , where

$$\beta = \frac{8\pi p_0}{B_0^2}, \quad (30)$$

and that the equilibrium quantities in the dense plasma are proportional to powers of r :

$$\mathbf{B}_0(r) = \bar{B} x^{-s} \hat{\theta}, \quad (31)$$

$$p_0(r) = \frac{\beta \bar{B}^2 x^{-2s}}{8\pi}, \quad \text{and} \quad (32)$$

$$\rho_0(r) = \bar{\rho} x^{-q}, \quad (33)$$

where

$$x \equiv \frac{r}{r_1}, \quad (34)$$

and r_1 , $\bar{B}$, and $\bar{\rho}$, are constants. It is also assumed that

$$g = \bar{g} x, \quad (35)$$

where $\bar{g}$ is a constant. Substituting equations (31) through (35) into equation (12), one finds that

$$q = 2s + 2, \quad (36)$$

$$v_A \propto r, \quad (37)$$

$$a \propto r, \quad (38)$$

and

$$\frac{g}{r} = \frac{\alpha v_A^2}{r^2}, \quad (39)$$

where

$$\alpha = [(q-2)(1+\beta) - 2]/2. \quad (40)$$

It is assumed that α and q are of order unity.

Using equations (31) through (40) in equation (26), one finds that

$$F = \bar{\rho} H x^{-q}, \quad (41)$$

where

$$H = D_A - \frac{h^2}{D_s} + (2-q) \frac{a^2}{r^2} \left(1 + \frac{mh}{D_s} \right) + \frac{v_A^2 - a^2}{r^2}. \quad (42)$$

Since D_A , D_s , h , and H are constant within the dense plasma, equation (25) has simple power-law solutions,

$$u_{1r} = x^b, \quad (43)$$

where

$$b^2 + b(2-q) + \Gamma = 0, \quad (44)$$

and

$$\Gamma = H \left(\frac{v_A^2}{r^2} + \frac{a^2 \omega^2}{r^2 D_s} \right)^{-1}. \quad (45)$$

4. DISPERSION RELATION FOR $k_z = 0$ NORMAL MODES IN THE CONSTANT- β EQUILIBRIUM

The dense plasma is assumed to have an inner edge at $r = r_1$, inside of which there is perfectly conducting plasma of vanishingly small density:

$$\rho \rightarrow 0 \quad \{\text{for } r < r_1\}. \quad (46)$$

Three different cases are considered for the behavior at large r : (1) the dense plasma has an outer edge at $r = r_2$, outside of which is non-magnetized plasma of vanishingly small density, (2) there is an outer edge at $r = r_2$, outside of which is low-density plasma with an axial magnetic field, and (3) the constant- β equilibrium profile is taken to extend to infinitely large r . All three cases lead to the essentially the same dispersion relation and stability criterion. In this section, cases (1) and (2) are treated; case (3) is discussed in appendix A.

For uniform rotation, the displacement ξ of a fluid element relative to where it would have been in the equilibrium flow is given by

$$\xi = \frac{i\mathbf{u}}{\omega}. \quad (47)$$

The boundary conditions at each of the boundaries between the high and low density plasmas are

$$||\hat{n} \cdot \xi|| = 0, \quad (48)$$

and

$$\left\| p + \frac{B^2}{8\pi} \right\| = 0, \quad (49)$$

where $\hat{n}$ is the unit normal to the perturbed boundary, and $||f||$ denotes the jump in f across the perturbed boundary. The Lagrangian perturbation of a quantity f , denoted δf , is related to the Eulerian perturbation, f_1 , by the equation $\delta f = f_1 + \xi \cdot \nabla f_0$. Equation (49) indicates that the Lagrangian total-pressure perturbation (magnetic plus thermal) is constant across the sharp boundary.

Since

$$\rho \rightarrow 0 \quad \{\text{for } r < r_1 \text{ or } r > r_2\}, \quad (50)$$

a and/or v_A must be arbitrarily large in the low-density plasma in order to maintain pressure balance in the equilibrium state. It is assumed that ω is fixed by the dynamics inside the dense plasma, so that high-frequency Alfvén and sound waves in the low-density plasma are not treated. For the comparatively low-frequency modes that are considered, the low-density plasma instantly rearranges itself to avoid the local compressions that would lead to thermal or magnetic-pressure perturbations. Within the low-density plasma, the Lagrangian total-pressure perturbation thus vanishes, as shown in appendix B. Equation (49) then implies that the Lagrangian total-pressure perturbation in the dense plasma also vanishes at the sharp boundaries:

$$p_1 + \xi \cdot \nabla p_0 + \frac{B_0 B_{1\theta}}{4\pi} + \xi \cdot \nabla \frac{B_0^2}{8\pi} = 0 \quad \{\text{for } r = r_1 \text{ or } r = r_2\}. \quad (51)$$

Using equations (19) and (22), one can rewrite equation (51) as

$$A_1 u_{1r} + A_2 \left(x \frac{du_{1r}}{dx} \right) = 0 \quad \left\{ \text{for } x = 1 \text{ or } x = \frac{r_2}{r_1} \right\}, \quad (52)$$

where

$$A_1 = \frac{a^2}{r^2} (D_s + mh), \quad (53)$$

and

$$A_2 = \frac{\omega^2 a^2 + D_s v_A^2}{r^2}. \quad (54)$$

The general solution to equation (25) for the constant- β equilibrium is a linear combination of the solutions given in equation (43),

$$u_{1r} = c_1 x^{b_1} + c_2 x^{b_2}, \quad (55)$$

where

$$b_1 = \left(\frac{q}{2} - 1 \right) + \sqrt{\left(\frac{q}{2} - 1 \right)^2 - \Gamma}, \quad (56)$$

and

$$b_2 = \left(\frac{q}{2} - 1 \right) - \sqrt{\left(\frac{q}{2} - 1 \right)^2 - \Gamma}. \quad (57)$$

Applying equation (52) at $x = 1$ and at

$$x_2 \equiv \frac{r_2}{r_1} \quad (58)$$

yields the matrix equation

$$\begin{pmatrix} A_1 + A_2 b_1 & A_1 + A_2 b_2 \\ A_1 + A_2 b_1 & x_2^{b_2 - b_1} [A_1 + A_2 b_2] \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (59)$$

Non-trivial solutions require that

$$(A_1 + A_2 b_1)(A_1 + A_2 b_2)(x_2^{b_2 - b_1} - 1) = 0. \quad (60)$$

One type of normal mode results from setting $x_2^{b_2 - b_1} = 1$, or, equivalently,

$$b_2 - b_1 = \frac{2\pi n i}{\ln(r_2/r_1)}, \quad n = \pm 1, \pm 2, \dots \quad (61)$$

Equations (56), (57) and (61) yield the dispersion relation

$$\Gamma - p_n = 0, \quad (62)$$

where

$$p_n = \left[\frac{\pi n}{\ln(r_2/r_1)} \right]^2 + \left(\frac{q}{2} - 1 \right)^2. \quad (63)$$

The Lagrangian total-pressure perturbation of these modes vanishes at $r = r_1$, $r = r_2$, and at $n - 1$ discrete values of r between r_1 and r_2 . Note that $n = 0$ in equation (61) does not correspond to an eigenmode, since it implies $b_1 = b_2$, $c_1 = -c_2$ from equation (59), and thus $u_{1r} = 0$.

A second type of normal mode results from having either

$$A_1 + A_2 b_1 = 0 \quad \text{and} \quad (64)$$

$$c_2 = 0, \quad (65)$$

or

$$A_1 + A_2 b_2 = 0 \quad \text{and} \quad (66)$$

$$c_1 = 0. \quad (67)$$

In either case, one sets $b = -A_1/A_2$ and substitutes into equation (44) to obtain

$$0 = \frac{a^4}{r^4} (D_s + mh)^2 + \left(\frac{v_A^2 D_s}{r^2} + \frac{a^2 \omega^2}{r^2} \right) \left[D_A D_s - h^2 + \frac{D_s (v_A^2 - a^2)}{r^2} \right]. \quad (68)$$

The Lagrangian total-pressure perturbation of these modes vanishes everywhere—they are the gravest modes of the system for any given m . The right-hand side of equation (68) is proportional to D_s , but the $D_s = 0$ roots are spurious; in deriving equation (25) it was assumed that $D_s \neq 0$, and it can be verified

that $D_s = 0$ does not lead to normal-mode solutions satisfying the boundary conditions.

Equations (62) and (68) can be rewritten in the form

$$0 = D_e D_R + S, \quad (69)$$

where

$$D_e = \omega^2 - G, \quad (70)$$

$$D_R = \omega^2 - \omega \frac{4mg\Omega}{rG} + \frac{m^2}{G} \left(\frac{g^2}{r^2} - \frac{\zeta a^2 v_A^2}{r^4} \right), \quad (71)$$

$$G = 4\Omega^2 + \frac{\zeta v_A^2}{r^2}, \quad (72)$$

and

$$\zeta = \begin{cases} p_n + m^2 - 1 & \text{for the roots of equation (62)} \\ m^2 - 1 & \text{for the roots of equation (68)} \end{cases}. \quad (73)$$

The value of S is different for equation (62) and equation (68). For equation (62)

$$S = \frac{4mg\Omega\omega^3}{rG} - \frac{m^2\omega^2}{G} \left(\frac{g^2}{r^2} - \frac{\zeta a^2 v_A^2}{r^4} \right) - \frac{a^2}{r^2} \left[\omega^2(q + \zeta) + m \left(2\omega\Omega - \frac{mg}{r} \right) \right], \quad (74)$$

whereas for equation (68)

$$S = -\frac{a^2 v_A^2}{r^2(a^2 + v_A^2)} \left[\omega^2 \left(1 - \frac{m^2 a^2}{v_A^2} \right) + \frac{m^2(m^2 - 1)a^2}{r^2} + 2m \left(2\omega\Omega - \frac{mg}{r} \right) \right] + \frac{4mg\Omega\omega^3}{rG} - \frac{m^2\omega^2(4\Omega^2 a^2 + g^2)}{Gr^2}. \quad (75)$$

It is now assumed that the dense plasma satisfies the low- β condition

$$a \ll v_A, \quad (76)$$

and is supported against gravity primarily by rotation, which implies that

$$v_A \ll v_{\text{rot}}, \quad (77)$$

where

$$v_{\text{rot}} \equiv |\Omega r|. \quad (78)$$

Equations (76) and (77) imply that S is negligible compared to the dominant terms in $D_e D_R$ for all values of ω , m , and p_n . The approximate roots of equation (69) are then obtained by setting $D_e D_R = 0$. When $D_e = 0$,

$$\omega \simeq \pm \sqrt{4\Omega^2 + \zeta v_A^2 / r^2}. \quad (79)$$

When $D_R = 0$,

$$\omega \simeq \frac{2mg\Omega r^2 \pm m v_A \sqrt{\zeta(4\Omega^2 a^2 r^2 - g^2 r^2 + \zeta a^2 v_A^2)}}{4\Omega^2 r^3 + \zeta r v_A^2}. \quad (80)$$

The errors in equations (79) and (80) vanish as $(a/v_A) \rightarrow 0$ and $(v_A/v_{\text{rot}}) \rightarrow 0$. Comparisons between equation (80) and numerical solutions of equations (62) and (68) are given in figure 3.

When a/v_A and v_A/v_{rot} are small (just how small will be seen below in the numerical solutions), both the approximate roots given in equations (79) and (80) and the exact roots of equations (62) and (68) are real for all m and n if and only if

$$|g| < 2|\Omega|a. \quad (81)$$

[To be precise, the exact stability criterion is slightly different from equation (81) due to small corrections associated with

nonzero a/v_A and v_A/v_{rot} .] It can be seen directly from equation (80) that the approximate roots are real for all m and n if and only if equation (81) is satisfied; that the exact roots are also real can be seen from the following argument. Let

$$D \equiv D_e D_R + S. \quad (82)$$

The four roots of the exact dispersion relation $D = 0$ are real when the plot of $D(\omega)$ passes through zero four times, as in figure 4. When $a \ll v_A$ and $v_A \ll v_{\text{rot}}$, D always has two local minima at $\omega \simeq \pm \sqrt{G/2}$ at which $D \simeq -G^2/4$. Also, $D \rightarrow +\infty$ as $|\omega| \rightarrow +\infty$. The question of stability turns upon whether D is non-negative for some value(s) of ω in the interval $(-\sqrt{G/2}, \sqrt{G/2})$. When $a \ll v_A$ and $v_A \ll v_{\text{rot}}$ and equation (81) is satisfied, the local maximum of $D_e D_R$ at $\omega = \omega_{\text{max}} \simeq 2mg\Omega/rG \in (-\sqrt{G/2}, \sqrt{G/2})$ is positive. Moreover, $S(\omega_{\text{max}})$ is negligible compared to the value of $D_e D_R$ at ω_{max} (unless the plasma is extremely close to marginal stability). Hence, the value of D is also positive at $\omega_{\text{max}} \in (-\sqrt{G/2}, \sqrt{G/2})$, and the four exact roots are real. On the other hand, when equation (81) is not satisfied, the local maximum of $D_e D_R$ at $\omega = \omega_{\text{max}}$ is negative, $D_e D_R$ is negative for all ω in the interval $(-\sqrt{G/2}, \sqrt{G/2})$, $|S| \ll |D_e D_R|$ for all ω in the interval $(-\sqrt{G/2}, \sqrt{G/2})$, $D_e D_R + S$ is negative for all ω in the interval $(-\sqrt{G/2}, \sqrt{G/2})$, and two of the exact roots have nonzero imaginary parts.

The $|m| = 1$ roots of equation (68) are a special case, since equation (80) gives a double root at $\omega = g/2\Omega r$. In this case, the local maximum of $D_e D_R$ occurs at $\omega = g/2\Omega r$, and $D_e D_R = 0$ at the local maximum. It is thus not clear whether the local maximum of $D_e D_R + S$ has a positive or negative value. In appendix C, a new approximate factorization of equation (68) is given for $|m| = 1$ and it is shown that the $|m| = 1$ roots are always stable when $a \ll v_A$ and $v_A \ll v_{\text{rot}}$, whether or not equation (81) is satisfied.

Figure 5 summarizes a stability analysis based upon numerical solutions of equations (62) and (68) for values of a/v_A and v_A/v_{rot} ranging from 0.01 to 1. A combination of parameters (a/v_A) and (v_A/v_{rot}) is said to be stable if the imaginary part of each of the numerically determined roots is less than 10^{-10} times the real part of the same root for all values of m from 1 to 100, and in the case of the roots to equation (62), for all values of n from 1 to 500, where $\ln(r_2/r_1)$ is taken to be 5, and q is taken to be 5. The roots are determined using an eigenvalue method (Press et al 1992) at double precision. Equation (81) describes the stability of the exact roots quite accurately provided $v_A/v_{\text{rot}} \lesssim 0.5$. For larger values of v_A/v_{rot} , equation (81) underestimates stability.

For any given values of a , Ω , and g , the most unstable mode has zero Lagrangian total-pressure perturbation everywhere and $\zeta = m^2 - 1$. Upon setting $d\omega_I/dm = 0$, where ω_I is the imaginary part of ω , one can show using equation (80) that the maximum growth rate γ_{max} occurs at

$$m_{\text{max}} \simeq \pm \frac{v_{\text{rot}}}{v_A} \sqrt{-6 + \sqrt{4 + \frac{8g^2}{\Omega^2 a^2}}} \quad (83)$$

and is given by

$$\gamma_{\text{max}} \simeq \frac{g}{v_A} f(\Omega^2 a^2 / g^2), \quad (84)$$

where f is a function that increases monotonically from 0 to 1 as its argument $\Omega^2 a^2 / g^2$ is decreased from 0.25 (marginal stability) towards 0. Because of equation (39), $\gamma_{\text{max}} \sim v_A/r$

when the plasma is not very near marginal stability. When $|g| > 2|\Omega|a$, modes with $\zeta = m^2 - 1$ as well as modes with $\zeta = p_n + m^2 - 1$ are unstable only when $\zeta < \zeta_{\text{crit}}$, where

$$\zeta_{\text{crit}} = \frac{r^2(g^2 - 4\Omega^2 a^2)}{a^2 v_A^2}. \quad (85)$$

5. DISCUSSION OF STABILITY CRITERION AND EIGENMODES

In the absence of background flow, the stability of an equilibrium to a small-amplitude displacement $\xi(\mathbf{x})$ is determined by whether $\xi(\mathbf{x})$ leads to an increase or decrease in δW , where δW is the sum of the perturbed magnetic, internal, and gravitational potential energies of the plasma (Bernstein et al 1958). If δW is positive, then the mode is stable—if the displacement increases with time initially, the initial bulk-motion kinetic energy present in the mode is converted into these other types of energy, and the plasma proceeds to oscillate stably. If δW is negative, then the magnetic, internal, and gravitational energies of the plasma can be tapped to drive an instability. In an equilibrium with flow, the kinetic energy of the flow is altered by plasma displacements, and stability is no longer determined solely by the increments to the magnetic, internal, and gravitational potential energies. The perturbed kinetic energy depends upon both the displacement $\xi(\mathbf{x})$ and its time derivative $\partial \xi / \partial t$, which gives rise to the phenomenon of overstability, in which both the real and imaginary parts of ω are nonzero, and which makes the stability analysis more complicated than in the static case (Frieman & Rotenberg 1960).

In this section, the physical basis of the stability criterion is explained using the properties of the eigenmodes. For any set of plasma parameters, the most unstable $k_z = 0$ modes are those described by equation (68) [as opposed to those described by equation (62)], for which the Lagrangian total-pressure perturbation vanishes everywhere. It is these modes that will be discussed in this section. The dispersion relation given by equation (68) has two branches, described approximately by equations (79) and (80). Equation (79) is an epicyclic-Alfvénic branch. When $m \sim O(1)$, one finds that $\omega = \pm 2\Omega$, $u_{1\theta} \simeq \mp i u_{1r}$, and $b \simeq -(\pm m + 1)a^2/v_A^2$. These circularly polarized epicyclic modes, which involve primarily a balance between inertia and the Coriolis force, are stable for all plasma parameters: when a fluid element in the dense plasma of figure 2 tries to sink towards $r = 0$, the Coriolis force pushes it in the $+\hat{\theta}$ direction (for positive Ω); as it moves in the $+\hat{\theta}$ direction, the Coriolis force pushes it back towards larger r . For $m^2 \gg v_{\text{rot}}^2/v_A^2$, the modes are stable Alfvén waves, with $\omega = \pm m v_A/r$, $|u_{1\theta}| \ll |u_{1r}|$, and $b \simeq -a^2/v_A^2$. The dominant force across field lines for these modes is magnetic tension.

Equation (80) is the Rayleigh-Taylor/sound-wave branch of the dispersion relation. It is shown in appendix D that the eigenmodes corresponding to equation (80) satisfy $|u_{1\theta}| \gg |u_{1r}|$, which indicates that these modes involve comparatively little field-line bending. For $m \sim O(1)$, $\omega \ll \Omega$. In the frame rotating with the plasma, the momentum equation at first order in perturbation amplitude can be written $-i\omega \rho_0 \mathbf{u}_1 = -2\rho_0 \Omega \hat{z} \times \mathbf{u}_1 + \dots$, where only the Coriolis force $\mathbf{F}_C$ has been written explicitly on the right-hand side. Since $\omega \ll \Omega$ and $\mathbf{u}_1$ is not along $\hat{z}$, some force must approximately cancel the Coriolis force. For $m \sim O(1)$, the near cancellation is provided by the gravitational force $\mathbf{F}_g$. Upon defining $F_{g,\parallel} = \mathbf{F}_g \cdot \mathbf{B}/B_0$ and $F_{C,\parallel} = \mathbf{F}_C \cdot \mathbf{B}/B_0$,

and using equation (18) to eliminate B_{1r} , one obtains

$$F_{g,\parallel} = \frac{\rho_0 m g u_{1r}}{\omega r}, \quad \text{and} \quad (86)$$

$$F_{C,\parallel} = -2\rho_0 \Omega u_{1r}. \quad (87)$$

It is shown in appendix D that $\rho_1 \simeq \rho_0 m u_{1\theta} / \omega r$ (that is, the density perturbation is dominated by the compression associated with $u_{1\theta}$), so that

$$\hat{r} \cdot \mathbf{F}_g \simeq -\frac{\rho_0 m g u_{1\theta}}{\omega r}, \quad (88)$$

while

$$\hat{r} \cdot \mathbf{F}_C = 2\rho_0 \Omega u_{1\theta}. \quad (89)$$

The phase velocity of a low- m mode in the $\hat{\theta}$ direction, $v_{\text{phase}} = \omega r / m$, is set by the requirement that $\mathbf{F}_C$ and $\mathbf{F}_g$ approximately cancel, which yields $v_{\text{phase}} \simeq g/2\Omega$, or $\omega \simeq mg/2\Omega r$. This phase velocity is in the eastwards direction when $g > 0$ —i.e., in the direction of rotation ($\omega/m\Omega > 0$). Because of the additional forces in the plasma, which are much smaller than either $\mathbf{F}_g$ or $\mathbf{F}_C$, the exact phase velocity will be either a little larger than $|g/2\Omega|$, in which case $|F_g| < |F_C|$, or a little smaller than $|g/2\Omega|$, in which case $|F_g| > |F_C|$.

It will prove useful to consider the radial momentum equation in somewhat greater detail. It is shown in appendix D that $|b| \ll 1$ for the eigenmodes of equation (80) with $m \sim O(1)$. This is related to the fact that the Lagrangian perturbation of the magnetic pressure, which is $\propto du_{1r}/dr$, is cancelled by the Lagrangian perturbation of the much smaller thermal pressure. (Although $|u_{1\theta}/u_{1r}|$ is large, it is not large enough to enable $|b|$ to be $\gtrsim 1$.) With the use of the relation $|b| \ll 1$, the vanishing of the Lagrangian total-pressure perturbation, equations (18) and (19), and the equation $\rho_1 = -\rho_0 \nabla \cdot \xi - \xi \cdot \nabla \rho_0$ (Frieman & Rotenberg 1960), the radial component of the momentum equation can be written

$$-i\omega \rho_0 u_{1r} \simeq \frac{\rho_0 u_{1\theta}}{\omega} \left(2\Omega\omega - \frac{mg}{r} \right) - \frac{i u_{1r} \rho_0 (m^2 - 1) v_A^2}{\omega r^2}. \quad (90)$$

The $(\rho_0 u_{1\theta}/\omega)(2\Omega\omega - mg/r)$ term is the radial component of what will be called the net gravitational force, $\mathbf{F}_g + \mathbf{F}_C$. The $-i u_{1r} \rho_0 (m^2 - 1) v_A^2 / \omega r^2$ term is the effective magnetic tension, which acts to restore a radially displaced fluid element in the dense plasma to its initial position. In order for a mode with $\omega \simeq mg/2\Omega r$ to propagate, the radial components of the effective tension and the net gravitational force must approximately cancel; otherwise, solving for ω yields $|\omega| \gtrsim v_A/r$, which is $\gg mg/2\Omega r$ for $m \sim O(1)$.

Because gravity and the Coriolis force nearly cancel, the low- m eigenmodes of equation (80) do not enjoy the full stabilizing effect that the Coriolis force has on epicyclic modes, nor the full destabilizing effect that gravity has on Rayleigh-Taylor modes in a static plasma. The physics behind the stability criterion can be understood by considering a mode with $m \sim O(1)$ and $|v_{\text{phase}}|$ slightly less than $|g/2\Omega|$, which implies that $|F_g|$ is slightly bigger than $|F_C|$ and that the net gravitational force acts in the same direction as gravity. If $|v_{\text{phase}}| < a$, then the dense plasma reacts in a way that is similar to how it would react if v_{phase} were 0: the maximum density enhancements occur at those points towards which the parallel component of the net gravitational force converges—the troughs of the perturbed magnetic field lines, as in figure 6. The induced pressure force opposes the parallel component of the net gravitational force. In the radial direction, the net gravitational force and the effective magnetic tension oppose each other and can therefore approximately cancel, as required for a mode with $\omega \simeq mg/2\Omega r$. On

the other hand, if $|\mathbf{v}_{\text{phase}}| > a$, then sound waves can not propagate information about the net gravity upstream of the wave. At any instant in time, fluid elements at the troughs of the perturbed field lines can not tell that the net gravitational force is trying to compress them, and the plasma reacts inertially to the parallel component of the net gravitational force. Plasma is accelerated towards positive $\hat{\theta}$ at those points where $B_{1r}/B_0 < 0$, and the resulting azimuthal compression causes the density enhancements to occur at the crests of the field lines when ω is real, as in figure 6. [This point can also be seen with the aid of equation (D18) in appendix D.] In this case, the net gravitational force in the radial direction acts in the same direction as the effective magnetic tension, providing much more radial force than is consistent with $\omega \simeq mg/2\Omega r$, and no stable mode at approximately that frequency exists.

Overstable modes, however, do exist. One way of thinking about this is the following. Suppose a low- m mode were to start propagating with a phase velocity greater than a and peak density enhancements at the crests of waves (as would be the case for real ω). The radial component of the net gravitational force and magnetic tension would then provide much more force than required to return the displaced plasma to its initial position. A fluid element displaced a distance ξ_{r0} towards $+\hat{r}$ would in response pass through its initial position and then move a distance $> \xi_{r0}$ towards $-\hat{r}$, leading to an oscillation of growing amplitude. If $\omega_I \neq 0$, where ω_I and ω_R are the imaginary and real parts of ω , respectively, then the phase relation between ρ_1 and u_{1r} is not as depicted in figure 6. When $m \sim \mathcal{O}(1)$, $|\omega_R| \gg |\omega_I|$ as in equation (80). $F_{g,\parallel}$ and $F_{C,\parallel}$ are then effectively sine waves of equal amplitude that are nearly 180° out of phase, and their sum is about 90° out of phase with each of them. The locations at which ρ_1 obtains its peak magnitude are then approximately the steepest points on the perturbed magnetic field lines, in phase with B_{1r} and u_{1r} . The peak magnitude of the radial component of the net gravitational force is then about 90° out of phase with ρ_1 and coincides with the troughs and crests of field lines. The radial component of the net gravity is then approximately cancelled by the effective magnetic tension.

When $|g| > 2|\Omega|a$, modes are stabilized when ζ is increased above ζ_{crit} because pressure begins to dominate over gravity in the dynamics along the magnetic field lines, changing the modes from Rayleigh-Taylor instabilities into something akin to sound waves. It is shown in appendix D that $|F_{g,\parallel}| > |F_{p,\parallel}|$ when $\zeta < \zeta_{\text{crit}}$ and $|g| > 2|\Omega|a$, where $F_{p,\parallel}$ is the component of the pressure force along the perturbed magnetic field lines in the dense plasma. It is also shown that as ζ is increased towards ζ_{crit} , $|F_{p,\parallel}|/|F_{g,\parallel}| \rightarrow 1$. For $\zeta > \zeta_{\text{crit}}$, $|F_{p,\parallel}| > |F_{g,\parallel}|$ for all m with one exception: $\zeta_{\text{crit}} < \zeta < g^2 r^2 / a^2 v_A^2$ on the branch of the dispersion relation with $\pm\Omega < 0$. This exception corresponds to the segment of the plot of ω_R in the upper-left panel of figure 3 just to the right of ζ_{crit} where the curve bifurcates, between the bifurcation point and where the plot of ω_R intersects $\omega_R = 0$; the plot of ω_R bifurcates at $m = 49$, and the exception corresponds to the smaller value of ω at $m = 50$.

To obtain further insight into the stability criterion, it is helpful to consider some limiting cases. Of course, if there is neither rotation nor magnetic field, the equilibrium of figure 2 is unstable to Rayleigh-Taylor modes as in the classic calculation of a dense fluid suspended above low-density fluid in slab geometry (Rayleigh 1883). If the orthogonal magnetic fields of figure 2 are present but the plasma isn't rotating, the situation is similar to the stationary plane-parallel problem solved by Gratton,

Gratton, & Gonzalez (1988) and Gonzalez & Gratton (1990). These authors discussed a number of cases, including that of a plasma at $y > 0$ suspended above vacuum at $y < 0$ with gravity pointing in the $-\hat{y}$ direction, all equilibrium quantities being functions of y alone, the magnetic field in the plasma pointing in the x direction, and the magnetic field in the vacuum pointing in the z direction. They found that the orthogonal magnetic fields act to stabilize the plasma, since the interface at $y = 0$ can not be perturbed without bending either the magnetic field in the plasma or the magnetic field in the vacuum. Compressibility, however, acts to destabilize the plasma; if the plasma is strongly compressed along the magnetic field, the gravitational force acting on regions of enhanced density can overcome magnetic tension. When the wave vector $\mathbf{k}$ of the perturbation in the $x-z$ plane is oriented along the magnetic field in the plasma at $y > 0$, magnetic tension on its own is not sufficient to stabilize small-scale modes: as $a \rightarrow 0$, the maximum unstable wavenumber approaches ∞ . Since thermal pressure limits the compressibility of the plasma along the magnetic field, magnetic tension and pressure together stabilize modes with $k^2 > k_{\text{crit}}^2$, where $k_{\text{crit}}^2 = g^2(v_A^2 + a^2)/a^2 v_A^4$. It can be shown that the condition $|k| > k_{\text{crit}}$ corresponds to pressure dominating over gravity in the dynamics along the perturbed magnetic field lines, in analogy to the stabilization of the modes described by equation (80) when $\zeta > \zeta_{\text{crit}}$. On the other hand, sufficiently large-wavelength modes ($|k| < k_{\text{crit}}$) are always unstable in the magnetized slab, whereas all $k_z = 0$ modes are stable in the rotating magnetized cylinder when $|g| < 2|\Omega|a$.

If there is rotation but no magnetic field, the plasma has a set of normal modes analogous to those described by equation (68) (i.e., $k_z = 0$ and zero Lagrangian pressure perturbation everywhere), with frequencies

$$\omega = \Omega \pm \sqrt{\Omega^2 - \frac{mg}{r}} \quad (91)$$

and

$$\omega = -\Omega \pm \sqrt{\Omega^2 + \frac{mg}{r}}. \quad (92)$$

These modes are circularly polarized ($u_{1\theta} = \pm u_{1r}$), stable at long wavelengths ($|m| < \Omega^2 r/g$), unstable when $m > \Omega^2 r/g$ in equation (91), and unstable when $m < -\Omega^2 r/g$ in equation (92). As $|m| \rightarrow \infty$, the growth time of unstable modes ($\sim 1/\sqrt{mg/r}$) is much shorter than the rotation period, and rotation can be neglected. The introduction of an azimuthal magnetic field into the dense plasma has a stabilizing influence insofar as magnetic tension eliminates these large- $|m|$ instabilities with $|u_{1\theta}| = |u_{1r}|$.

The magnetic field in the equilibrium of figure 2, however, can also be destabilizing: as the field strength is increased, the plasma becomes increasingly supported by magnetic pressure, thereby increasing the effective gravity g . With the use of equation (39), the stability criterion can be rewritten as an upper limit on the Alfvén speed:

$$v_A^2 < \left(\frac{2}{\alpha}\right) a v_{\text{rot}} \quad (93)$$

(where a , v_A , and v_{rot} are evaluated in the dense plasma). In the low- β limit, pressure balance requires that the magnetic-field strength B_{vert} in the low-density plasma at $r < r_1$ be approximately equal to the field strength in the dense plasma near $r = r_1$, $\bar{B}$. Assuming a dense plasma of pure hydrogen with $q = 5$, $\gamma = 5/3$, $\beta \ll 1$, $v_{\text{rot}} = 150$ km/s [from the Galactic rotation curve at $r = 150$ pc (Ruzmaikin & Shukurov 1981)], and

$T = 70$ K and $n = 10^4 \text{ cm}^{-3}$ near $r = r_1$, one can rewrite equation (93) as

$$B_{\text{vert}} < 1.1 \text{ mG}. \quad (94)$$

Although it is tempting to infer from equation (94) that mG vertical magnetic fields at the Galactic center can be stably confined by a ring of horizontal magnetic fields threading molecular material, further study is needed to extend the present analysis to more realistic equilibria.

6. OTHER TYPES OF INSTABILITIES

Sections 2 through 5 treated the global stability of the equilibrium pictured in figure 2 to $k_z = 0$ modes. This section addresses three other classes of instabilities: (1) local interchange and non-axisymmetric quasi-interchange modes in a uniformly rotating plasma with azimuthal magnetic field, (2) global instabilities in the 1D equilibrium of figure 2 with $k_z \neq 0$, and (3) quasi-interchange and magneto-rotational instabilities in a two-dimensional differentially rotating equilibrium such as the one pictured in figure 1.

Local modes in a plasma with an azimuthal magnetic field that rotates uniformly about the z axis have been treated by a number of authors. For the constant- β equilibrium profile of section 3, local axisymmetric interchange instabilities are suppressed primarily because of the strongly stabilizing specific-angular-momentum profile when $a \ll v_A \ll v_{\text{rot}}$ (Newcomb 1962, Acheson & Gibbons 1978, Rogers & Sonnerup 1986, Ferriere et al 1999). Such modes are further stabilized since the specific entropy and magnetic flux per unit mass of an azimuthal flux loop B/pr are taken to increase with r in the constant- β equilibrium of section 4. The results of Ferriere et al (1999) on a wide class of one-dimensional equilibria can be used to show that local non-axisymmetric quasi-interchange modes in a uniformly rotating plasma with azimuthal magnetic field are stable when two conditions are met assuming $a \ll v_A \ll v_{\text{rot}}$: (1) the “magnetic Brunt-Väisälä” frequency ω_0 must be real, and (2) $|g| < 2|\Omega|a$ (to lowest order in a/v_A and v_A/v_{rot}). For the constant- β equilibrium of section 4, $\omega_0^2 \simeq qg/2r$ and the first condition is met. The second condition is the same as equation (81), and thus when the $k_z = 0$ modes are stable, so are the quasi-interchange modes. The quasi-interchange modes have $\omega \simeq mg/2\Omega r$, and are stable (provided ω_0 is real) when their phase velocity along the magnetic field is less than the sound speed, just as for $k_z = 0$ modes with $m \sim O(1)$.

Global modes in the equilibrium of figure 2 with $k_z \neq 0$ and $m \sim O(1)$ are analogous to the external kink instability in static plasmas. An $m = 1$ mode is illustrated in figure 7 [figure 9.23 of Freidberg (1987)]. Because the strength of the azimuthal field increases where the field lines are compressed, the Lorentz force of the azimuthal field acts to increase the perturbation. On the other hand, the magnetic tension of the axial magnetic field opposes the perturbation. In the static reversed field pinch, in which $B_\theta \sim B_z$, the azimuthal magnetic field wins out over the axial field and the mode is unstable when the axial wavelength k_z^{-1} is greater than roughly the radius of the plasma cross section (Friedberg 1987). How this instability mechanism affects the stability criterion of the rotating equilibrium of figure 2 is left for future work.

Newcomb (1961) showed that the convective stability criterion for a compressible stratified magnetized slab when $k_{\parallel} = 0$ is less stringent than the stability criterion in the limit $k_{\parallel} \rightarrow 0$,

where $k_{\parallel}$ is the component of the wave vector along the magnetic field. This is because when $k_{\parallel} = 0$, Newcomb’s type 2 modes, which involve motion primarily along the magnetic field, become pure translations which are neutrally stable ($\omega = 0$). The $k_{\parallel} = 0$ stability criterion is thus the one that applies to Newcomb’s type 1 modes, which involve motion primarily across field lines. On the other hand, when $k_{\parallel}$ is small but nonzero, the type 2 modes allow destabilizing compressions of the plasma along the magnetic field and can be unstable even when type 1 modes are stable. The distinction between $k_{\parallel} = 0$ and $k_{\parallel} \rightarrow 0$ in Newcomb’s work is not directly analogous to the distinction between $k_z = 0$ and $k_z \rightarrow 0$ in the equilibrium of figure 2, in that $k_{\parallel}$ corresponds to m since it is the dense plasma that dominates the dynamics of the unstable modes.

The two-dimensional equilibrium of figure 1 is subject to instabilities not present in the one-dimensional infinite-cylinder equilibrium. If the vertical (along z) stratification does not satisfy the Schwarzschild criterion for a static non-magnetized gas, then the plasma is unstable to local non-axisymmetric quasi-interchange modes with $k_r \gg k_z$ and $k_r r/m \gg 1$ (Terquem & Papaloizou 1996). The displacements in such modes are essentially parallel to the spin axis, and thus the Coriolis force vanishes, preventing the type of rotational stabilization that occurs for quasi-interchange modes in the one-dimensional infinite-cylinder equilibrium. If there is differential rotation with $d|\Omega|/dr < 0$, the plasma is subject to the magnetorotational instability (MRI) (Ogilvie & Pringle 1996, Terquem & Papaloizou 1996, Balbus & Hawley 1998). Global three-dimensional numerical simulations suggest that turbulence excited by the MRI will be present quite generally in astrophysical disks (Hawley, Balbus, & Stone 2001). It is thus interesting to ask how the stability of the sharp boundary in figure 1 is affected by differential rotation in a quiescent equilibrium flow as well as by differential rotation with turbulence. These questions are left for future work.

7. ESTIMATES OF FIELD STRENGTHS IN RADIO FILAMENTS

Lang et al. (1999b) found that if the magnetic energy density in the Northern Thread is equal to the relativistic-particle energy density, and if the relativistic protons and electrons have equal energies, then the field strength is $\simeq 140 \mu\text{G}$. Comparable equipartition field strengths have been found for the Radio Arc (Tsuboi et al. 1986, Yusef-Zadeh & Morris 1987b, Reich 1994). However, the equipartition estimate may lead to an underestimate of the field strength, particularly since the geometry of the filaments allows relativistic particles to stream directly out of the Galactic-center region.

All of the radio filaments that have been studied in detail near the Galactic center appear to be interacting with at least one randomly moving molecular cloud, and yet most are undeformed at the sites of interaction (Morris & Serabyn 1996). If the magnetic field were dynamically insignificant, one would expect the turbulent motions of the clouds to cause turbulent bulk motions in the hot plasma that would deform the frozen-in magnetic field lines. The lack of deformation gives a lower limit on the field strength in the filaments that depends upon how many independently moving clouds (or clumps within a cloud) interact with a single filament.

If a vertical filament interacts with a single molecular cloud at a single point along its length, then the lower limit on its field strength is fairly small. Since the vertical field B_{vert} threads the very tenuous hot plasma, there is very little mass to anchor

the ends of the vertical field line above and below the Galactic disk. Any deformation in the field line in the galactic plane by an interaction with a molecular cloud is carried away along the field line at the Alfvén speed of the hot medium, $v_{A,hot}$. In order for the visible deformations in the filaments to be small, the Alfvén speed in the hot medium must be much larger than the turbulent speed of the clouds v_{cloud} with respect to the ends of the filament. The angle at which a field line is bent will be $\sim v_{cloud}/v_{A,hot}$, as depicted in figure 8. The condition $v_{A,hot} \gg v_{cloud}$ can be written

$$B_{vert} \gg n_{hot}^{1/2} \left(\frac{v_{cloud}}{2 \text{ km/s}} \right) \mu\text{G}, \quad (95)$$

where n_{hot} is the number density of the hot plasma in cm^{-3} . If $v_{cloud} = 30 \text{ km/s}$, and $n_{hot} = 0.3$, then $B_{vert} \gg 8 \mu\text{G}$.

As pointed out by Yusef-Zadeh & Morris (1987a) and Morris & Serabyn (1996), if a filament interacts with clouds at more than one point along its length, as happens in the Radio Arc (Morris 2001), then the lower limit on the field strength may be considerably greater than in equation (95). For certain cloud shapes, a filament could conceivably avoid the appearance of deformation by rapidly slipping around the moving clouds. For some cloud shapes, however, this would not be possible, and to avoid deformation a filament would have to dynamically oppose the force applied by the clouds. If this requires that the magnetic pressure in the filament exceed the ram pressure in the clouds, then $B_{vert} > 1 \text{ mG}$ (Yusef-Zadeh & Morris 1987a). If a single cloud consists of independently moving clumps, then a similar argument would apply to a filament interacting with a single cloud (Morris & Serabyn 1996).

8. CONCLUSION

Because of the simplifying assumptions made in this paper, further work is needed to determine the extent to which the stability criterion given by equation (81) applies to more realistic equilibria, and to extend the analysis to modes with $k_z \neq 0$. If equation (81) does describe, even approximately, the stability of the boundary between the high and low-density plasmas in systems with differential rotation, a finite extent along the z axis, and more general radial profiles—say by evaluating the terms in

equation (81) near the inner edge of the dense plasma—then it appears that a pervasive mG vertical magnetic field can be confined at the Galactic center. Equation (81) would then also be relevant to other accreting systems with strong central poloidal magnetic fields.

A pervasive mG vertical magnetic field in the inner 150 pc of the Galaxy would have important implications for the origin of the Galactic magnetic field. If the vertical field originated from radial inflow over the lifetime of the Galaxy, then the strength of the average vertical magnetic field in the Galactic disk at the time of Galaxy formation, B_i , can be estimated by magnetic-flux conservation. If the average mass accretion rate into the Galactic-center region is $0.3 M_\odot/\text{yr}$ (Morris & Serabyn 1996), and if the surface density of gas in the disk has remained constant over the lifetime of the Galaxy, then all the gas that was within roughly 10 kpc of the Galactic center at the time of Galaxy formation has been accreted into the Galactic-center region. If the magnetic field was frozen in to the interstellar plasma, then the vertical magnetic flux that initially threaded the central 10 kpc of the disk, $\Phi_i = B_i \pi (10 \text{ kpc})^2$, is now concentrated near the Galactic center. If there is a pervasive vertical magnetic field at the Galactic center today, it is unlikely that it contains regions of oppositely directed vertical fields, since the parallel geometry of field lines would permit rapid destruction of the oppositely directed fields by turbulent diffusion and reconnection. The current magnetic flux through the Galactic-center region corresponding to such a pervasive field, Φ_f , can thus be estimated by multiplying field strength times area. If there is a mG vertical field throughout the central 150 pc of the Galaxy, then $\Phi_f = \pi (150 \text{ pc})^2 \text{ mG}$. Setting $\Phi_i = \Phi_f$ yields $B_i \sim 10^{-7} \text{ G}$ (Chandran et al 2000), which would be consistent with a turbulent dynamo operating in the protogalaxy (Kulsrud et al. 1997).

I thank Amitava Bhattacharjee, Eric Blackman, Steve Cowley, Ken Gayley, Cornelia Lang, and Mark Morris for valuable discussions. This work was supported by the National Science Foundation under grant AST 00-98086 and by the Department of Energy under grant DE-FG02-01ER54658 and grant DE-FC02-01ER54651.

APPENDIX

DISPERSION RELATION WHEN CONSTANT- β AZIMUTHAL-FIELD EQUILIBRIUM EXTENDS FROM $r = r_1$ TO $r = \infty$

In this case, the boundary conditions are that the Lagrangian total-pressure perturbation vanishes at $r = r_1$ and that the behavior as $r \rightarrow \infty$ is such that the kinetic energy of the mode is finite. As in section 4, the boundary conditions can be satisfied in two ways. The first is for both c_1 and c_2 to be non-zero. The boundary condition at $r = r_1$ is then given by the first row of equation (59), and a finite kinetic energy requires that the radicals in equations (56) and (57) be purely imaginary; if they are real or complex, then the r^{b_1} term causes the kinetic energy to diverge, where $\sqrt{\dots}$ is taken to be the root with a positive real part. The resulting dispersion relation is the same as equation (62), but with p_n replaced by $[(0.5q - 1)]^2 + N$, where N is any positive real number. The second way is to have $u_{1r} \propto r^b$ with $b = (0.5q - 1) - \sqrt{(0.5q - 1)^2 - 1}$, which leads to equation (68) and corresponds to a mode in which the Lagrangian total-pressure perturbation vanishes everywhere.

DIFFERENTIAL EQUATION FOR $k_z = 0$ NORMAL MODES IN THE LOW-DENSITY AXIAL-FIELD REGION

The differential equation for $k_z = 0$ normal modes in a uniformly rotating plasma with axial magnetic field $[\mathbf{B}_0 = B_0(r)\hat{z}]$ is

$$\frac{d}{dr} \left[\frac{\rho_0(a^2 + v_A^2)}{D_{ms}r} \frac{d}{dr} (ru_{1r}) \right] + rKu_{1r}, \quad (B1)$$

where

$$D_{ms} = \omega^2 - \frac{m^2(a^2 + v_A^2)}{r^2}, \quad (B2)$$

$$K = -\frac{d}{dr} \left[\frac{m\rho_0 M(a^2 + v_A^2)}{\omega^2 r^2 D_{ms}} \right] + \frac{\rho_0}{r} - \frac{\rho_0}{\omega^2} \frac{d}{dr} \left(\frac{g}{r} \right) - \frac{M^2 \rho_0}{\omega^2 r D_{ms}}, \quad (\text{B3})$$

and

$$M = \frac{mg}{r} - 2\Omega\omega. \quad (\text{B4})$$

When $g^* = 0$ (i.e., $g = -\Omega^2 r$), equation (B1) reduces to the constant- Ω limit of equation (3a) of Spies (1978). In the limit $\rho_0 \rightarrow 0$, with ρ_0 constant, equation (B1) reduces to

$$\frac{d}{dr} \left[r \frac{d}{dr} (ru_{1r}) \right] + m^2 u_{1r} = 0, \quad (\text{B5})$$

with solutions

$$u_{1r} = r^l, \quad (\text{B6})$$

with

$$l = -1 \pm m. \quad (\text{B7})$$

Writing the Lagrangian pressure perturbation as δp and the Lagrangian perturbation of B_z as δB_z , one finds that

$$\delta p + \frac{B_0 \delta B_z}{4\pi} = \frac{irB_0^2 \omega(2 + \gamma\beta)}{8\pi m^2 (a^2 + v_A^2)} \left[\frac{d}{dr} (ru_{1r}) + \frac{mMu_{1r}}{\omega^2} \right], \quad (\text{B8})$$

which vanishes as $\rho \rightarrow 0$ for fixed ω , where ω is set by the dynamics inside the dense plasma.

APPROXIMATE FACTORIZATION OF DISPERSION RELATION [EQUATION (68)] FOR $|m| = 1$.

For the $|m| = 1$ roots of equation (68), the dispersion relation can be written as

$$D_e \tilde{D}_R + S = 0, \quad (\text{C1})$$

where $\tilde{D}_R = (\omega - \omega_+)(\omega - \omega_-)$, and

$$\omega_{\pm} = \frac{1}{2\Omega} \left\{ \frac{g}{r} + \frac{a^2}{r^2} \left[-1 \pm \left(1 - \frac{g^2}{4a^2\Omega^2} \right) \right] \right\}. \quad (\text{C2})$$

The value of S can be recalculated and is different from the values in equations (74) and (75). As in the case $|m| \neq 1$, when $a \ll v_A$ and $v_A \ll v_{\text{rot}}$, $D_e \tilde{D}_R + S$ has local minima at $\omega \simeq \pm \sqrt{G/2}$ at which $D_e \tilde{D}_R + S \simeq -G^2/4$, and stability again turns upon whether $D_e \tilde{D}_R + S$ becomes non-negative in the interval $\omega \in (-\sqrt{G/2}, \sqrt{G/2})$. The value of $D_e \tilde{D}_R$ has a local maximum at $\omega = (\omega_+ + \omega_-)/2$ at which $D_e \tilde{D}_R$ is always positive. It can be shown that $S \ll D_e \tilde{D}_R$ at $\omega = (\omega_+ + \omega_-)/2$, and thus $D_e \tilde{D}_R + S$ is always positive at $\omega = (\omega_+ + \omega_-)/2$, which guarantees that the four exact roots are always real.

PROPERTIES OF THE EIGENMODES

This appendix treats the eigenmodes corresponding to equation (68), for which the Lagrangian total-pressure perturbation vanishes everywhere. All of the results of this section assume that

$$a \ll v_A \ll v_{\text{rot}}, \quad (\text{D1})$$

and a number of the equations to be presented assume that the plasma is not very near marginal stability, i.e.,

$$\left| \frac{g^2 - 4\Omega^2 a^2}{g^2 + 4\Omega^2 a^2} \right| \gg \frac{a}{v_A}, \quad \text{and} \quad \left| \frac{g^2 - 4\Omega^2 a^2}{g^2 + 4\Omega^2 a^2} \right| \gg \frac{v_A}{v_{\text{rot}}}. \quad (\text{D2})$$

Upon substituting equation (80) into equation (24), one finds that

$$D_s \simeq -\frac{m^2}{r^2 G^2} \left[2\Omega \sqrt{a^2 G - g^2} \mp \frac{g v_A \sqrt{\zeta}}{r} \right]^2, \quad (\text{D3})$$

where the $\simeq$ sign has the same meaning as in equation (80), that the error in the equation vanishes as $a/v_A \rightarrow 0$ and $v_A/v_{\text{rot}} \rightarrow 0$. When ω is real, $(a^2 G - g^2) > 0$, D_s is real, and $D_s < 0$. This implies that all stable modes on the Rayleigh-Taylor branch of the dispersion relation [equation (80)] travel more slowly than sound waves. Upon substituting equation (80) into equation (23), one finds that

$$h \simeq \pm \frac{m v_A \sqrt{\zeta}}{r^2 G} \left(2\Omega \sqrt{a^2 G - g^2} \mp \frac{g v_A \sqrt{\zeta}}{r} \right) + \frac{m a^2}{r^2}. \quad (\text{D4})$$

Given equation (D2), the ma^2/r^2 term on the right-hand side is negligible compared to the other term for $|m| > 1$. It can be shown that

$$\left| \frac{h}{D_s} \right| \gg 1 \quad (\text{D5})$$

for $|m| > 1$.

When

$$1 < m \ll v_{\text{rot}}/a \quad (\text{D6})$$

and equation (D2) is satisfied, it can be shown that $|\omega^2 a^2| \ll |D_s v_A^2|$,

$$b \simeq -\frac{a^2 m h}{v_A^2 D_s}, \quad (\text{D7})$$

and, for unstable modes with $a^2 G - g^2 < 0$,

$$|b| \simeq \frac{a^2}{v_A} \sqrt{\frac{\zeta G}{g^2 - 4\Omega^2 a^2}}. \quad (\text{D8})$$

The value of $|b|$ in equation (D8) is $\ll 1$ when $|m| \sim \mathcal{O}(1)$ and increases monotonically with increasing ζ , reaching a maximum value of

$$|b|_{\text{max}} \simeq \frac{gr}{v_A^2} = \alpha \quad (\text{D9})$$

at ζ_{crit} , the largest value of ζ for which modes are still unstable [equation (85)]. When $m \sim \mathcal{O}(1)$, $|b| \ll 1$ for stable modes as well.

Given equations (D2) and (D6), it can be shown that $|(ma^2/r)du_{1r}/dr| \ll |hu_{1r}|$ and

$$u_{1\theta} \simeq -\frac{ihu_{1r}}{D_s}. \quad (\text{D10})$$

Equation (D5) then implies that

$$|u_{1\theta}| \gg |u_{1r}|. \quad (\text{D11})$$

Equations (D7) and (D10) can be used to show that

$$\nabla \cdot \mathbf{u}_1 \simeq \frac{imu_{1\theta}}{r}. \quad (\text{D12})$$

Since $|\nabla \cdot \mathbf{u}_1| \gg |u_{1r}/r|$, the relation (Frieman & Rotenberg 1960)

$$p_1 = -\gamma p_0 \nabla \cdot \xi - \xi \cdot \nabla p_0 \quad (\text{D13})$$

can be used to show that

$$p_1 \simeq -\gamma p_0 \nabla \cdot \xi \simeq -\frac{ia^2 \rho_0 m h u_{1r}}{\omega r D_s}. \quad (\text{D14})$$

Similarly, the relation $\rho_1 = -\rho_0 \nabla \cdot \xi - \xi \cdot \nabla \rho_0$ (Frieman & Rotenberg 1960) can be used to show that

$$\rho_1 \simeq \frac{\rho_0 m u_{1\theta}}{\omega r} \simeq -\frac{i \rho_0 m h u_{1r}}{\omega r D_s}. \quad (\text{D15})$$

If one takes the dot product of equation (3) with $\mathbf{B}/B_0$, uses equation (4) to eliminate $\mathbf{B}_1$ and equation (D13) to eliminate p_1 , one finds that

$$-i\omega \rho_0 u_{1\theta} = -\frac{u_{1r} \rho_0}{\omega} \left(2\Omega\omega - \frac{mg}{r} \right) - \frac{ma^2 \rho_0 \nabla \cdot \mathbf{u}_1}{\omega r}. \quad (\text{D16})$$

The term on the left-hand side represents inertia in the frame of reference rotating with the plasma, the terms $F_{C,\parallel} = -u_{1r} 2\Omega \rho_0$, $F_{g,\parallel} = u_{1r} mg \rho_0 / \omega r$, and $F_{p,\parallel} = -ma^2 \rho_0 (\nabla \cdot \mathbf{u}_1) / \omega r$ are the components of the Coriolis, gravitational, and pressure forces along the perturbed magnetic field line in roughly the $+\hat{\theta}$ direction. One can show that for modes with complex ω ,

$$\left| \frac{F_{p,\parallel}}{F_{g,\parallel}} \right|^2 \simeq \frac{a^2 G \zeta}{g^2 \zeta_{\text{crit}}} \quad (\text{D17})$$

which is less than 1 for $\zeta < \zeta_{\text{crit}}$, but which approaches 1 as $\zeta \rightarrow \zeta_{\text{crit}}$ (as ζ is increased towards ζ_{crit} , $a^2 G$ is increased towards g^2). The stabilization of modes as ζ is increased to ζ_{crit} thus coincides with pressure becoming equal to gravity in the dynamics along field lines. With the use of equation (D12), equation (D16) can be rewritten as

$$-iD_s u_{1\theta} \simeq -u_{1r} \left(2\Omega\omega - \frac{mg}{r} \right). \quad (\text{D18})$$

Equations (D15) and (D18) show that the sign of D_s is critical for determining the phase of ρ_1 relative to u_{1r} . Let $F\hat{b}$ be the sum of the components of the gravitational and Coriolis forces along the magnetic field. As explained in section 5, if ω is real and $D_s < 0$, the peaks in the perturbed density will occur at the points of convergence of $F\hat{b}$. If ω is real and $D_s > 0$, the peaks in the perturbed density will occur at the points of divergence of $F\hat{b}$.

REFERENCES

- Acheson, D. J., and Gibbons, M. P. 1978, *J. Fluid Mech.*, 85, 743
- Balbus, S. A., & Hawley, J. F. 1998, *Rev. Mod. Phys.*, 70, 1
- Bally, J., Stark, A., Wilson, R., and Henkel, C. 1988, *ApJ*, 324, 223
- Benford, G. 1988, *ApJ*, 333, 735
- Bernstein, I., Frieman, E., Kruskal, M., & Kulsrud, R. 1958, *Proc. Roy. Soc. (London)*, A244, 17
- Chandran, B. D. G., Cowley, S. C., & Morris, M. 2000, *ApJ*, 528, 723
- Chandrasekhar, S. 1961, *Hydrodynamic and hydromagnetic stability* (New York: Dover)
- Chudnovsky, E. M., Field, G. B., Spergel, D. N., & Vilenkin, A. 1986, *Phys. Rev. D*, 34, 944
- Ferriere, K. M., Zimmer, C., & Blanc M. 1999, *J. Geophys. Res.*, 104, 17335
- Friedberg, J. 1987, *Ideal Magnetohydrodynamics* (New York: Plenum; p. 425)
- Frieman, E., and Rotenberg, M. 1960, *Rev. Mod. Phys.*, 32, 898
- Gilman, P. 1970, *ApJ*, 162, 1019
- Gonzalez, A., and Gratton, J. 1990, *Plasm. Phys. Contr. Nuc. Fus.*, 32, 3
- Gratton, J., Gratton, F., Gonzalez, A. 1988, *Plasma Phys. Contr. Nuc. Fus.*, 30, 435
- Hawley, J., Balbus, S., & Stone, J. 2001, *ApJ*, 554L, 49
- Heyvaerts, J., Norman, C., & Pudritz, R. E. 1988, *ApJ*, 330, 718
- Koyama, K., Maeda, Y., Sonobe, T., Takeshima, T., Tanaka, Y., & Yamauchi, S. 1996, *Publ. Astron. Soc. Japan*, 48, 249
- Kulsrud, R. M., Cen, R., Ostriker, J. P., Ryu, D. 1997, *ApJ*, 480, 481
- Lang, C., Anantharamaiah, K., Kassim, N., & Lazio, T. 1999, 521, L41
- Lang, C., Morris, M., & Echevarria, L. 1999, *ApJ*, 526, 727
- Lesch, H., & Reich, W. 1992, *A&A*, 264, 493
- Lubow, S. H., Papaloizou, J. C. B., & Pringle, J. E. 1994, *MNRAS*, 267, 235
- Morris, M. 1996, in *Unsolved Problems of the Milky Way*, eds. L. Blitz and P. Teuben (Netherlands: IAU, p. 247)
- Morris, M. 2001, private communication
- Morris, M., and Serabyn, E. 1996, *Ann. Rev. Astron. Astrophys.*, 34, 645
- Newcomb, W. 1961, *Phys. Fluids*, 4, 391
- Newcomb, W. 1962, *Nucl. Fusion*, suppl., part 2, 451
- Press, W., Teukolsky, S., Vetterling, W., & Flannery, B. 1992, *Numerical Recipes in C* (Cambridge: Cambridge University Press)
- Rayleigh, Lord 1883, *Proc. Lond. Math. Soc.*, 14, 170
- Reich, W. 1994, in *Nuclei of Normal Galaxies: Lessons from the Galactic Center*, eds. Genzel, R., Harris, A. I. (Dordrecht: Kluwer, p. 55)
- Rogers, B., Sonnerup, B. 1986, *J. Geophys. Res.*, 91, 8837
- Rosso, F., & Pelletier, G. 1993, *A&A*, 270, 416
- Ruzmaikin, A., & Shukurov, M. 1981, *Sov. Astron.*, 25, 553
- Shore, S. N., & LaRosa, T. N. 1999, *ApJ*, 521, 587
- Sofue, Y., & Fujimoto, M. 1987, *Publ. Astron. Soc. Jpn.*, 39, 843
- Spies, G. O. 1978, *Phys. Fluids*, 21, 580
- Talwar, S. P. 1960, *J. Fluid Mech.*, 9, 581
- Terquem, C., & Papaloizou, J. 1996, *MNRAS*, 279, 767
- Tsuboi, M., Inoue, M., Handa, T., Tabara, H., Kato, T. et al. 1986, *Astron. J.*, 92, 818
- Tsuboi, M., Kawabata, T., Kasuga, T., Handa, T., Kato, T. 1995, *Publ. Astron. Soc. Jpn.*, 47, 829
- Yamauchi, S., Kawada, M., Koyama, K., Kunieda, H., & Tawara, Y. 1990, *ApJ*, 365, 532
- Yusef-Zadeh, F., and Morris, M. 1987a, *Astron. J.*, 94, 1178
- Yusef-Zadeh, F., and Morris, M. 1987b, *ApJ*, 322, 721




FIG. 1.— A simplified model for an equilibrium of strong vertical fields at the Galactic center.




FIG. 2.— The simplified one-dimensional equilibrium that is analyzed in this paper.

TABLE 1
NOTATION

Symbol	Equation where introduced	Meaning
$\rho, p, \mathbf{u}, \mathbf{B}$	(2), (3)	mass density, flow velocity, pressure, and magnetic field; (0 indicates equilibrium value, 1 indicates perturbed quantity)
g^*, g	(3), (11)	true gravity, effective gravity
Ω	(8)	uniform angular velocity
a, v_A, v_{rot}	(13), (14), (78)	sound speed, Alfvén speed, rotation speed
γ	(13)	adiabatic index
m, k_z, n	(15), (61)	azimuthal, axial, and radial mode numbers
σ, ω	(15), (17)	frequency in inertial frame, frequency in rotating frame
h, F, H, Γ, p_n	(23), (26), (42), (45), (63)	convenient abbreviations
D_s, D_A, D_e, D_R	(24), (27), (70), (71)	dispersion-relation polynomials for the sound wave, Alfvén wave, epicyclic-Alfvén mode, and Rayleigh-Taylor mode
β	(30)	ratio of thermal to magnetic pressure
x	(34)	normalized radius
q	(33)	exponent in constant- β equilibrium ($\rho_0 \propto r^{-q}$)
α	(39), (40)	order-unity factor in constant- β equilibrium
b	(43)	exponent of r in power-law solutions for u_{1r}
ξ	(47)	Lagrangian displacement
ζ	(73)	measure of the wavenumber
ζ_{crit}	(85)	maximum value of ζ corresponding to an unstable mode
$\mathbf{F}_g, \mathbf{F}_C, \mathbf{F}_p$	(86), (87), (D17)	perturbed gravitational, Coriolis, and pressure forces
B_{vert}	(94)	strength of the pervasive vertical field at the Galactic center




FIG. 3.— Comparison of approximate analytic solution to dispersion relation for Rayleigh-Taylor modes [equation (80)] and numerical solutions of equations (62) and (68). The solid (dashed) lines represent the real (imaginary) part of ω/Ω from equation (80), while the crosses (diamonds) give the real (imaginary) part of numerical solutions of equations (62) and (68). The upper two panels correspond to equation (68), and the lower two panels correspond to equation (62). In the upper-right panel, $a/v_A = 0.1$ and $v_A/v_{\text{rot}} = 0.1$, and the plasma is stable. In the other three panels, $a/v_A = 0.01$ and $v_A/v_{\text{rot}} = 0.2$, and the plasma is unstable. For all of the plots, $q = 5$.




FIG. 4.— Plots of $D_e D_R$ (solid line) and $D_e D_R + S$ (dashed line), with S given by equation (74), $a/v_A = 0.2$, $v_A/v_{\text{rot}} = 0.2$, $m = 10$, $n = 30$, $\ln(r_2/r_1) = 5$, and $q = 5$.




FIG. 5.— The crosses indicate combinations of parameters (evaluated in the dense plasma) for which the numerical solutions of equations (62) and (68) indicate instability. The dots indicate parameters for which the numerical solutions are real. The solid line corresponds to the approximate analytic stability criterion equation (93).




FIG. 6.— Eigenmodes corresponding to equation (80) with $m \sim \mathcal{O}(1)$ and a Lagrangian total-pressure perturbation that vanishes everywhere. $F_{g,\parallel}$ and $F_{C,\parallel}$ are the components of the gravitational and Coriolis forces along the perturbed magnetic field lines in the dense plasma. When ω is real and $v_{\text{phase}} < a$, the plasma is compressed at the points towards which $F_{g,\parallel} + F_{C,\parallel}$ converges, so that the perturbed pressure force opposes $F_{g,\parallel} + F_{C,\parallel}$. When ω is real and $v_{\text{phase}} > a$, the plasma responds inertially and is compressed at the points away from which $F_{g,\parallel} + F_{C,\parallel}$ diverges. In the latter case, magnetic tension is unable to balance the perturbed radial gravitational force, and no stable mode exists.




FIG. 7.— An $m = 1$ external kink mode.




FIG. 8.— Because the field lines in the hot low-density plasma are not anchored above or below the Galactic plane, they remain almost straight if they interact with a cloud at only a single point along their length provided the Alfvén speed in the hot plasma $v_{A, \text{hot}}$ is $\gg v_{\text{cloud}}$.